

Prairie Fen Companion *Plant facts*

Poison sumach

Toxicodendron vernix

C=10

Color: green

5 regular parts




Plant type: Flowering shrub

Bloom period: June

Distinguishing characteristics: Alternate compound leaves have pointed, shiny leaflets and a red stem. The species has clusters of small green flowers in June that ripen to white berries that stay on the plant through the growing season, sometimes into winter. This is a woody plant that grows to 15 ft tall. Bark is a dull grey. This species produces the same oil in its sap as poison ivy and oak. Contact with leaves and stems (even in winter) will result in a severe rash if you are sensitive to this plant.

Habitat quality: This species often grows mixed with other shrubs, and is found both in low and high-quality fens. It is an indicator of wet conditions.

Where can you find this plant?



MICHIGAN STATE
UNIVERSITY



Michigan
Natural
Features
Inventory

The Nature
Conservancy
Protecting nature. Preserving life.™




The Stewardship Network

Developed by: Doug Landis and Anna Fiedler, MSU Department of Entomology. Funding support: National Fish and Wildlife Foundation, Lynn and Thelma MacCready Forest and Wildlife Endowment, MSU, and Hanes Trust of the Michigan Botanical Club. Partners: The Nature Conservancy, Michigan Natural Features Inventory, The Stewardship Network, Michigan DNR Landowner Incentive Program. For more information on native plants and prairie fens, go to www.nativeplants.msu.edu.

Where can you find this plant?



Indiana



New York



Ohio



Wisconsin

For more information on native plants and prairie fens, go to www.nativeplants.msu.edu

Key description

C=


Coefficient of conservatism. This is a value that ranges from 0 for non-native invasives to 10 for plants that would only be expected to be found in undisturbed, high quality plant communities. It is a general guideline for whether the plant would be likely to be found in an intact prairie fen (not filled in with shrubs and without invasive species). However, C values may be high for some species that are not found in prairie fen but would be part of another wetland such as a marsh. They are included here because they are a widely accepted measure of habitat quality in the Midwest (<http://1.usa.gov/FQAMethod>).

Flower type

Classifications here follow those of Newcomb's wildflower guide (<http://amzn.com/0316604429>).

Number of regular parts

The flower has this number of petals or petal-like parts that are symmetrical from the flower center (radial), with each similar to the other in shape, size, and color. There may be 3 to 7 regular parts. See image at right.



Aster

These flowers have regular parts and are symmetrical from the center, but there are more than 7. Asters have a set of disc flowers in the center of the flower and a set of ray flowers outside of the disc flowers, often called petals. They are one group (genus) within the family Asteraceae, and there are many species in this genus.

Flowers not readily obvious

The flowers of plants such as grasses, sedges, and cattail are not obvious and are often confused with the fruits (seeds) of these species. While this website does not include flower descriptions for these species, they do flower.

Irregular

The flower is not symmetrical from the center but is symmetrical down a line (bilateral). See image at right.



No flowers

A number of primitive plants, including ferns, do not flower but make spores in order to reproduce.

Parts indistinguishable

These species either have parts so small their number is difficult to determine or have no petal-like parts. This group includes goldenrods, other species with small individual flowers, and plants in the family Asteraceae that have more than 7 parts, but the parts do not form distinguishable, symmetrical ray flowers (which are often called petals).