

# grapes.msu.edu

[Home](#)[About Us](#)[Search](#)[Newsletters](#)[Weather /  
Climate](#)[Viticulture](#)[Pest  
management](#)[Scouting guide](#)[Publications](#)[Calendar of  
events](#)[Industry links](#)[Contacts](#)

Download [Adobe Acrobat Reader](#) to view pdf files.

## Diuron injury

Tom Zabadal , MSU Horticulture

[Home](#) > [Scouting guide](#)> diuron injury

Diuron herbicide injury causes leaf veins to appear yellow to cream-colored. Severe cases can cause stunting of vines. This injury is often associated with light soils or areas where soil has eroded so that the vine roots are near the soil surface. Vines may need more than one year to outgrow this injury.

### Additional information

- Search [MSU Extension News for Agriculture](#) site
- Search [MSU Fruit CAT Alert](#) newsletter for articles.
- [MSU Diagnostic Services](#)



Leaves showing the typical yellow veinal pattern of diuron herbicide injury.  
Photo: T. Zabadal

[Site map](#)[Copyright/Linking](#)

Funding for this web site provided by [Project GREEN](#), [American Farmland Trust](#), [EPA Region 5's Strategic Agricultural Initiative](#) program, [The National Foundation for IPM Education](#), the [Center for Agricultural Partnerships](#) and the [MSU Integrated Pest Management Program](#) in collaboration with [MSU Extension](#) and the [Michigan Agricultural Experiment Station](#). Partially support from [NC-IPM Center](#).

05/24/11 Contact: [E. Haney](#)